

Parity proofs of the Kochen-Specker theorem based on the Lie algebra E8

Mordecai Waegell and P.K. Aravind

May 5, 2015

Abstract The 240 root vectors of the Lie algebra E8 lead to a system of 120 rays in a real 8-dimensional Hilbert space that contains a large number of parity proofs of the Kochen-Specker theorem. After introducing the rays in a triacontagonal representation due to Coxeter, we present their Kochen-Specker diagram in the form of a “basis table” showing all 2025 bases (i.e., sets of eight mutually orthogonal rays) formed by the rays. Only a few of the bases are actually listed, but simple rules are given, based on the symmetries of E8, for obtaining all the other bases from the ones shown. The basis table is an object of great interest because all the parity proofs of E8 can be exhibited as subsets of it. We show how the triacontagonal representation of E8 facilitates the identification of substructures that are more easily searched for their parity proofs. We have found hundreds of different types of parity proofs, ranging from 9 bases (or contexts) at the low end to 35 bases at the high end, and involving projectors of various ranks and multiplicities. After giving an overview of the proofs we found, we present a few concrete examples of the proofs that illustrate both their generic features as well as some of their more unusual properties. In particular, we present a proof involving 34 rays and 9 bases that appears to provide the most compact proof of the KS theorem found to date in 8 dimensions.

1 Introduction

The exceptional Lie algebra E8 plays a role in a number of physical theories such as supergravity and heterotic string theory[1]. Here we show that its system of root vectors can be used to exhibit a large number of “parity proofs” of the Kochen-Specker (KS) theorem[2] ruling out the existence

M.Waegell^{1,2}, P.K.Aravind¹

¹Physics Department, Worcester Polytechnic Institute, Worcester, MA 01609, U.S.A.

²Institute for Quantum Studies, Chapman University, Orange, CA.

E-mail: caiw@wpi.edu, paravind@wpi.edu

of noncontextual hidden variables theories as viable alternatives to quantum mechanics. The fact that the root vectors of E8 could be made to serve this end was pointed out, in different ways, by Lisonek et. al.[3] and by Ruuge and van Oystaeyen[4]. However the proof in [4] is unrelated to parity proofs, while [3], though proving that 2^{1940} parity proofs exist, does not list even a single example of such a proof. The purpose of this paper is to supplement the observations in [3] and [4] by (a) presenting a general framework (namely, the “basis table”) within which the parity proofs can be exhibited, (b) showing how the symmetries of E8 can be exploited to simplify the search for its parity proofs, (c) providing an overview of the parity proofs found by our search and, finally, (d) presenting a few concrete examples of the proofs in order to convey some feeling for their variety and intricacy.

Parity proofs of the KS theorem are appealing because they take no more than simple counting to verify. What a parity proof is, and how it accomplishes its goals, are matters that will be explained later in this paper.

We make a few remarks about E8, to shed light on the way it is used in this paper. For us E8 is simply a collection of 240 vectors (namely, its roots) in a real 8-dimensional Euclidean space. These vectors define the vertices of a semiregular polytope discovered by Thorold Gosset[5] and sometimes described by the symbol 4_{21} . The vectors have eight real coordinates that can be chosen in a variety of ways. A particularly judicious choice, for our purposes, is the “triacontagonal representation” of Coxeter and Shepard[6]. In this representation the coordinates of the vectors are chosen in such a way that if only the first two coordinates are retained (which amounts to projecting the vectors orthogonally from eight dimensions down to two), the tips of the vectors lie at the vertices of eight regular triacontagons lying on concentric circles. Such a projection is shown in Fig.10 of [6]. The eight rings of thirty points are easily picked out in the figure, while the dense network of lines connecting pairs of points are projections of the 6720 edges of Gosset’s polytope.

Two slight modifications of this figure will convert it into the Kochen-Specker diagram of E8: the first is that only one member of each pair of diametrically opposite vertices should be retained, since one is concerned with rays rather than vectors in a proof of the KS theorem; and the second is that the line segments corresponding to the edges of Gosset’s polytope should be replaced by new segments that connect only pairs of vertices that correspond to orthogonal rays. Rather than construct such a diagram, we later present a “basis table” that conveys essentially the same information in a more useful form. The basis table is simply a listing of all the bases (i.e., sets of eight mutually orthogonal rays) in the system. Because there are 2025 bases, and this is too large a number to display explicitly, we list just a few bases and give a simple set of rules (based on the symmetries of E8) for generating all the bases from the ones shown. It should be pointed out that the entire basis table can in fact be generated from any one of its elements by applying products of all

possible powers of three basic symmetry operations of E8. The basis table is of central importance in this paper because all the parity proofs of E8 can be exhibited as subsets of it.

We say a few words about the broader area in which this work is set, to provide some perspective. After their initial discovery in two-qubit systems[8, 9, 10, 11, 12], parity proofs were discovered in three-[13, 14] and higher-qubit[15, 16] systems, in systems of rays derived from the four-dimensional regular polytopes [17, 18, 19] and the root systems of exceptional Lie algebras[20] and, very recently, in a remarkably compact six-dimensional system of complex rays[21] in connection with which an experiment has also been reported[22]. Aside from revealing the many guises in which quantum contextuality can arise in spaces of different dimensionality, parity proofs are interesting because they have a variety of applications: they can be used to derive state-independent inequalities for ruling out noncontextuality[23, ?, 25, 26, ?, 28] and Bell inequalities for identifying fully nonlocal correlations[29]; they have applications to quantum games[30], quantum zero-error communication[31], quantum error correction[32, 33] and the design of relational databases[34]; they can be used to witness the dimension of quantum systems[35]; and they underlie surprising phenomena such as the quantum pigeonhole effect[36, 37, 38]. Although the KS theorem is theoretically compelling, it has been argued[39, 40, 41] that the finite precision of real measurements nullifies practical attempts at verifying it. There is some debate about this matter[42], but it should be mentioned that methods of establishing contextuality that are not open to this objection have been proposed[43, 44, 45, 46]. Spekkens[47] has recently expanded upon the conditions that must be satisfied by realistic experiments that claim to rule out noncontextual ontological models. It has been argued in [48] that contextuality is the source of the speedup in many quantum information protocols. This brief survey is far from complete, but it serves to show that the KS theorem and quantum contextuality are at the heart of many current research efforts.

The plan of this paper is as follows. Section 2 introduces the triacontagonal representation of the 120 rays derived from the root vectors of E8, and shows how their symmetries can be exploited to give an efficient construction of their basis table. Section 3 points out some interesting substructures within E8 that have been shown in the past to give rise to proofs of the KS theorem. Section 4 reviews the notion of a parity proof and shows how the triacontagonal representation of E8 facilitates the identification of subsets of its bases, of distinct symmetry types, that each house a large number of parity proofs. The significance of these smaller subsets is that they are far more easily searched for parity proofs than the entire system. After giving an overview of the parity proofs we found among the different subsets of bases, we present a few examples of the parity proofs that illustrate their important features. Section 5 concludes with a discussion of our results.

2 The E8 system: rays and bases

The 240 root vectors of E8 come in 120 pairs, with the members of each pair being the negatives of each other. Choosing just one member from each pair yields the 120 rays associated with E8. Each ray has eight real coordinates, which may be chosen in a variety of ways. We use a coordinatization due to Richter[7], which differs from the one introduced earlier by Coxeter and Shepard[6]. Let $\omega = \exp(\frac{i\pi}{30})$ and let $\tau = \frac{1+\sqrt{5}}{2}$ be the golden ratio. Define a, b, c and d as the positive numbers satisfying the equations

$$\begin{aligned} 2a^2 &= 1 + 3^{-1/2}5^{-1/4}\tau^{3/2}, & 2b^2 &= 1 + 3^{-1/2}5^{-1/4}\tau^{-3/2} \\ 2c^2 &= 1 - 3^{-1/2}5^{-1/4}\tau^{-3/2}, & 2d^2 &= 1 - 3^{-1/2}5^{-1/4}\tau^{3/2}. \end{aligned}$$

For any integer n , let $c_n = \omega^n + \omega^{-n} = 2\cos(\frac{n\pi}{30})$ and define the quantities

$$\begin{aligned} r_1 &= a/c_9, & r_2 &= b/c_9, & r_3 &= c/c_9, & r_4 &= d/c_9, \\ r_5 &= a/c_3, & r_6 &= b/c_3, & r_7 &= c/c_3, & r_8 &= d/c_3. \end{aligned}$$

The 120 rays $|i\rangle$ ($i = 1, \dots, 120$) are then defined as

$$\begin{aligned} |n+1\rangle &= (r_1\omega^{2n}, r_4\omega^{22n}, r_6\omega^{14n+1}, r_7\omega^{26n+1}) && \text{for } 0 \leq n \leq 14 \\ |n+16\rangle &= (r_4\omega^{2n}, -r_1\omega^{22n}, r_7\omega^{14n+1}, -r_6\omega^{26n+1}) && \text{for } 0 \leq n \leq 14 \\ |n+23\rangle &= (r_7\omega^{29+2n}, -r_6\omega^{19+22n}, -r_1\omega^{24+14n}, r_4\omega^{18+26n}) && \text{for } 8 \leq n \leq 14 \\ |n+38\rangle &= (r_7\omega^{29+2n}, -r_6\omega^{19+22n}, -r_1\omega^{24+14n}, r_4\omega^{18+26n}) && \text{for } 0 \leq n \leq 7 \\ |n+38\rangle &= (r_6\omega^{29+2n}, r_7\omega^{19+22n}, r_4\omega^{24+14n}, r_1\omega^{18+26n}) && \text{for } 8 \leq n \leq 14 \\ |n+53\rangle &= (r_6\omega^{29+2n}, r_7\omega^{19+22n}, r_4\omega^{24+14n}, r_1\omega^{18+26n}) && \text{for } 0 \leq n \leq 7 \\ |n+61\rangle &= (r_8\omega^{2n}, -r_5\omega^{22n}, -r_3\omega^{14n+1}, r_2\omega^{26n+1}) && \text{for } 0 \leq n \leq 14 \\ |n+76\rangle &= (r_5\omega^{2n}, r_8\omega^{22n}, -r_2\omega^{14n+1}, -r_3\omega^{26n+1}) && \text{for } 0 \leq n \leq 14 \\ |n+83\rangle &= (r_2\omega^{29+2n}, r_3\omega^{19+22n}, -r_8\omega^{24+14n}, -r_5\omega^{18+26n}) && \text{for } 8 \leq n \leq 14 \\ |n+98\rangle &= (r_2\omega^{29+2n}, r_3\omega^{19+22n}, -r_8\omega^{24+14n}, -r_5\omega^{18+26n}) && \text{for } 0 \leq n \leq 7 \\ |n+98\rangle &= (r_3\omega^{29+2n}, -r_2\omega^{19+22n}, r_5\omega^{24+14n}, -r_8\omega^{18+26n}) && \text{for } 8 \leq n \leq 14 \\ |n+113\rangle &= (r_3\omega^{29+2n}, -r_2\omega^{19+22n}, r_5\omega^{24+14n}, -r_8\omega^{18+26n}) && \text{for } 0 \leq n \leq 7, \end{aligned}$$

with each ray being an 8-component column vector whose (real) components are given by the real and imaginary parts of the four complex numbers listed for it¹. We will use $\langle i|$ to denote the 8-component row vector that is the transpose of $|i\rangle$.

Let us denote by the letters A,...,H each consecutive set of 15 rays (thus A denotes rays 1-15, B rays 16-30, etc.). If we add to each group of 15 rays all their negatives, we get groups of 30 vectors whose first two coordinates define the vertices of regular triacontagons in the plane, with the triacontagons corresponding to the eight letter groups being concentric to one another. This

¹ For example, the components of the column vector $|3\rangle$, in the proper order, are $r_1 \cos(\frac{2\pi}{15}), r_1 \sin(\frac{2\pi}{15}), r_4 \cos(\frac{22\pi}{15}), r_4 \sin(\frac{22\pi}{15}), r_6 \cos(\frac{29\pi}{30}), r_6 \sin(\frac{29\pi}{30}), r_7 \cos(\frac{53\pi}{30}), r_7 \sin(\frac{53\pi}{30})$.

is just the triacontagonal representation of the roots of E8 (or of the vertices of Gosset's polytope 4_{21}) mentioned in the introduction. Although the coordinates we have introduced for the rays are identical to those of Richter[7], our numbering of the rays is a bit different from his (in essence, we have swapped some of his triacontagons and rotated some of them relative to the others for convenience in the presentation of some of our results).

A straightforward calculation shows that each of the 120 rays is orthogonal to 63 others and that the rays form 2025 bases. Each ray occurs in 135 bases and its only companions in these bases are the 63 other rays it is orthogonal to. We will denote this system of rays and bases by the symbol $120_{135}-2025_8$, with the subscript on the left indicating the multiplicity of each of the rays (i.e., the number of bases it occurs in) and that on the right the number of rays in each basis. The product of the numbers in the left half of the symbol equals the product on the right, as it should. The basis table of E8 (i.e., the complete list of all its bases) is *saturated*, by which we mean that all the orthogonalities between its rays are represented in it. Because of this, the basis table is completely equivalent to the Kochen-Specker diagram of its rays². However it has the great advantage over the Kochen-Specker diagram that it is easy to interpret and work with.

Later we will encounter other systems of rays and bases having a high degree of symmetry, and the notation we have introduced above is easily modified to deal with such cases. For example, a system of 45 rays that forms 15 bases, with 30 of the rays being of multiplicity 2 and the other 15 of multiplicity 4 can be represented by the symbol $30_2 15_4-15_8$ (again the sum of the products of each number on the left with its subscript equals the product of the number and its subscript on the right). The parity proofs we will present later, which are subsets of the basis table, can also be described by symbols of this kind.

We now present the basis table of E8. Figure 1 shows 15 bases that contain all 120 rays once each. The entire basis table can be derived from these 15 bases by permuting the rays in them in the manner we now describe. Let V be the permutation of order 9 with the cycle decomposition³ $V = (1\ 5\ 9\ 13\ 53\ 40\ 82\ 105\ 11)(2\ 91\ 90\ 55\ 28\ 42\ 54\ 119\ 49)(3\ 47\ 38\ 66\ 31\ 30\ 41\ 103\ 12)(4\ 10\ 51\ 89\ 117\ 106\ 87\ 27\ 36)(6\ 93\ 97\ 101\ 86\ 71\ 48\ 69\ 113)(7\ 14\ 79\ 67\ 33\ 29\ 64\ 32\ 100)(8\ 95\ 99\ 45\ 44\ 92\ 112\ 63\ 78)(15\ 104\ 34\ 46\ 109\ 77\ 118\ 107\ 85)(16\ 120\ 98\ 60\ 61\ 75\ 18\ 35\ 68)(17\ 73\ 20\ 24\ 59\ 96\ 58\ 57\ 94)(19\ 23\ 76\ 52\ 84\ 56\ 21\ 25\ 37)(22\ 74\ 116\ 108\ 115\ 72\ 62\ 50\ 70)(65\ 80\ 88\ 102\ 83\ 110\ 114\ 81\ 111)(26\ 43\ 39)$ and W the permutation of order 15 with the cycle decomposition $W = (1\ \dots\ 15)(16\ \dots\ 30)(31\ \dots\ 45)(46\ \dots\ 60)(61\ \dots\ 75)(76\ \dots\ 90)(91\ \dots\ 105)(106\ \dots\ 120)$, where the dots signify all the integers between the two extremes. Let each basis in

² This is a graph whose vertices correspond to the rays and whose edges join vertices corresponding to orthogonal rays

³ By the cycle $(1\ 5\ 9\ 13\ \dots\ 11)$, we mean the permutation in which 1 goes to 5, 5 to 9, 9 to 13 ... and 11 to 1.

Figure 1 be assigned the label $(0, 0, l)$, where the first two labels are fixed and the third varies, in integer steps, from 0 to 14. Then any other member of the basis table, which is assigned the label (n, m, l) with $0 \leq l, n \leq 14$ and $0 \leq m \leq 8$, can be generated by applying suitable powers of W and V to one of the bases in Fig.1, as described by the equation $(n, m, l) = W^n V^m (0, 0, l)$. The number of bases that can be generated in this way is $15 \cdot 9 \cdot 15 = 2025$, which is the entire basis table.

Basis label	Basis
(0,0,0)	1 7 62 66 70 73 107 111
(0,0,1)	29 115 33 11 74 61 5 52
(0,0,2)	64 34 85 102 19 88 101 94
(0,0,3)	58 78 92 42 47 110 112 116
(0,0,4)	55 117 80 51 96 106 41 108
(0,0,5)	27 81 84 82 21 59 114 14
(0,0,6)	10 23 38 103 37 56 53 113
(0,0,7)	2 72 44 104 95 32 48 25
(0,0,8)	17 89 9 76 54 26 119 77
(0,0,9)	87 109 4 16 22 86 79 49
(0,0,10)	100 31 120 45 60 57 13 30
(0,0,11)	91 90 43 15 75 20 105 63
(0,0,12)	99 71 36 28 93 39 6 50
(0,0,13)	24 46 67 35 118 12 3 69
(0,0,14)	83 8 40 97 98 18 65 68

Fig. 1 Fifteen bases of the E8 system, involving the rays 1-120 once each. The first column shows the three-index label of each basis.

In Figure 2 we show the 9 blocks of bases obtained by applying all powers of V to the block of Figure 1. The remaining blocks of bases can be obtained by applying powers of W to these nine blocks. Since an application of W amounts, for the most part, to increasing the ray numbers by one, these other blocks are easily written down.

The construction we have given of the basis table can be compressed even further by introducing a permutation of order 15, which we will term U , whose cycle decomposition is given by the eight columns of numbers obtained by aligning the bases in Fig.1 and reading down them vertically⁴. Then all the bases can be generated from the first basis of Fig.1 by applying powers of the permutations U , V and W to it, as described by the equation

$$(n, m, l) = W^n V^m U^l (0, 0, 0) \quad , \quad (1)$$

with $0 \leq l \leq 14$, $0 \leq m \leq 8$ and $0 \leq n \leq 14$. This procedure works even if an arbitrary basis is substituted for $(0, 0, 0)$ as the seed basis. The three-index

⁴ To be explicit, $U = (1 \ 29 \ 64 \ \dots \ 83)(7 \ 115 \ 34 \ \dots \ 8) \ \dots \ (111 \ 52 \ 94 \ \dots \ 68)$, where there are 8 cycles and each consists of 15 numbers.

1 7 62 66 70 73 107 111 29 115 33 11 74 61 5 52 64 34 85 102 19 88 101 94 58 78 92 42 47 110 112 116 55 117 80 51 96 106 41 108 27 81 84 82 21 59 114 14 10 23 38 103 37 56 53 113 2 72 44 104 95 32 48 25 17 89 9 76 54 26 119 77 87 109 4 16 22 86 79 49 100 31 120 45 60 57 13 30 91 90 43 15 75 20 105 63 99 71 36 28 93 39 6 50 24 46 67 35 118 12 3 69 83 8 40 97 98 18 65 68	5 14 50 31 22 20 85 65 64 72 29 1 116 75 9 84 32 46 15 83 23 102 86 17 57 8 112 54 38 114 63 108 28 106 88 89 58 87 103 115 36 111 56 105 25 96 81 79 51 76 66 12 19 21 40 6 91 62 92 34 99 100 69 37 73 117 13 52 119 43 49 118 27 77 10 120 74 71 67 2 7 30 98 44 61 94 53 41 90 55 39 104 18 24 11 78 45 48 4 42 97 26 93 70 59 109 33 68 107 3 47 113 110 95 82 101 60 35 80 16	9 79 70 30 74 24 15 80 32 62 64 5 108 18 13 56 100 109 104 110 76 83 71 73 94 95 63 119 66 81 78 115 42 87 102 117 57 27 12 72 4 65 21 11 37 58 111 67 89 52 31 3 23 25 82 93 90 50 112 46 45 7 113 19 20 106 53 84 49 39 2 107 36 118 51 98 116 48 33 91 14 41 60 92 75 17 40 103 55 28 26 34 35 59 1 8 44 69 10 54 101 43 97 22 96 77 29 16 85 47 38 6 114 99 105 86 61 68 88 120
13 67 22 41 116 59 104 88 100 50 32 9 115 35 53 21 7 77 34 114 52 110 48 20 17 99 78 49 31 111 8 72 54 27 83 106 94 36 3 62 10 80 25 1 19 57 65 33 117 84 30 47 76 37 105 97 55 70 63 109 44 14 6 23 24 87 40 56 2 26 91 85 4 107 89 60 108 69 29 90 79 103 61 112 18 73 82 12 28 42 43 46 68 96 5 95 92 113 51 119 86 39 101 74 58 118 64 120 15 38 66 93 81 45 11 71 75 16 102 98	53 33 74 103 108 96 34 102 7 70 100 13 72 68 40 25 14 118 46 81 84 114 69 24 73 45 8 2 30 65 95 62 119 36 110 87 17 4 47 50 51 88 37 5 23 94 80 29 106 56 41 38 52 19 11 101 28 22 78 77 92 79 93 76 59 27 82 21 91 43 90 15 10 85 117 61 115 113 64 55 67 12 75 63 35 20 105 3 42 54 39 109 16 58 9 99 112 6 89 49 71 26 86 116 57 107 32 98 104 66 31 97 111 44 1 48 18 120 83 60	40 29 116 12 115 58 46 83 14 22 7 53 62 16 82 37 79 107 109 111 56 81 113 59 20 44 95 91 41 80 99 50 49 4 114 27 73 10 38 70 89 102 19 9 76 17 88 64 87 21 103 66 84 23 1 86 42 74 8 118 112 67 97 52 96 36 105 25 90 39 55 104 51 15 106 75 72 6 32 28 33 3 18 78 68 24 11 47 54 119 26 77 120 57 13 45 63 93 117 2 48 43 71 108 94 85 100 60 34 31 30 101 65 92 5 69 35 98 110 61
82 64 108 3 72 57 109 110 79 74 14 40 50 120 105 19 67 85 77 65 21 111 6 96 24 92 99 90 103 88 45 70 2 10 81 36 20 51 66 22 117 83 23 13 52 73 102 32 27 25 12 31 56 76 5 71 54 116 95 107 63 33 101 84 58 4 11 37 55 26 28 34 89 104 87 18 62 93 100 42 29 47 35 8 16 59 1 38 119 49 43 118 98 94 53 44 78 97 106 91 69 39 48 115 17 15 7 61 46 30 41 86 80 112 9 113 68 60 114 75	105 32 115 47 62 94 77 114 67 116 79 82 70 98 11 23 33 15 118 80 25 65 93 58 59 112 45 55 12 102 44 22 91 51 111 4 24 89 31 74 106 110 76 53 84 20 83 100 36 37 3 30 21 52 9 48 119 108 99 85 78 29 86 56 57 10 1 19 28 43 42 46 117 34 27 35 50 97 7 54 64 38 68 95 120 96 5 66 49 2 39 107 60 17 40 92 8 101 87 90 113 26 69 72 73 104 14 75 109 41 103 71 88 63 13 6 16 61 81 18	11 100 72 38 50 17 118 81 33 108 67 105 22 60 1 76 29 104 107 88 37 80 97 57 96 63 44 28 3 83 92 74 90 89 65 10 59 117 30 116 87 114 52 40 56 24 110 7 4 19 47 41 25 84 13 69 49 115 45 15 8 64 71 21 94 51 5 23 42 39 54 109 106 46 36 68 70 101 14 119 32 66 16 99 98 58 9 31 2 91 26 85 61 73 82 112 95 86 27 55 6 43 113 62 20 34 79 18 77 103 12 48 102 78 53 93 120 75 111 35

Fig. 2 The nine blocks of bases obtained by applying all powers of V , from 0 to 8, to the block of Figure 1. The powers increase as one goes from left to right and up to down. Applying V to the block at the bottom right gives back the block at the top left, which is just the block of Fig.1.

label (n, m, l) serves as a convenient shorthand for the basis if one wishes to avoid listing all its rays.

Instead of describing U, V and W by the permutations they perform on the rays, one can represent them by the 8×8 orthogonal matrices

$$U = -|29\rangle\langle 1| - |115\rangle\langle 7| - |33\rangle\langle 62| + |11\rangle\langle 66| + |74\rangle\langle 70| + |61\rangle\langle 73| - |5\rangle\langle 107| - |52\rangle\langle 111| \quad (2)$$

$$V = |5\rangle\langle 1| + |14\rangle\langle 7| - |50\rangle\langle 62| - |31\rangle\langle 66| - |22\rangle\langle 70| - |20\rangle\langle 73| - |85\rangle\langle 107| + |65\rangle\langle 111| \quad (3)$$

and

$$W = |2\rangle\langle 1| + |8\rangle\langle 7| + |63\rangle\langle 62| + |67\rangle\langle 66| + |71\rangle\langle 70| + |74\rangle\langle 73| + |108\rangle\langle 107| + |112\rangle\langle 111| , \quad (4)$$

as one can easily check by applying them to the column vectors representing the rays and verifying that they produce the desired permutations.

We can construct all the symmetries of E8 by looking at the mappings of a fixed basis on to all the bases of the system. Let $x_1x_2 \cdots x_8$ and $y_1y_2 \cdots y_8$ be two bases and let $y'_1y'_2 \cdots y'_8$ be some permutation of the numbers in the latter. Consider the orthogonal transformation

$$T = (-1)^{n_1}|y'_1\rangle\langle x_1| + (-1)^{n_2}|y'_2\rangle\langle x_2| + \cdots + (-1)^{n_8}|y'_8\rangle\langle x_8| , \quad (5)$$

where $n_i \in (0, 1)$ for $1 \leq i \leq 8$. Keeping the basis $x_1x_2 \cdots x_8$ fixed and letting $y_1y_2 \cdots y_8$ vary, the number of transformations of the form (5) that one can construct is the number of possibilities for the variable basis (2025) times the number of permutations of the variable basis labels (8!) times the number of possibilities for the signs of the terms (2^8). However an investigation shows that only 1/30 of the 8! permutations of the basis labels lead to symmetries of E8, so the total number of its symmetries is $2025 \cdot \frac{8!}{30} \cdot 2^8 = 696729600$, which equals the number of $192 \cdot 10!$ given by Coxeter[5] as the order of the symmetry group of Gosset's polytope 4_{21} .

3 Substructures within the E8 system

The E8 system contains a number of interesting substructures that yield proofs of the KS theorem. These substructures have all been studied in the past, and we discuss each of them briefly.

Two interesting substructures are the Lie algebras E7 and E6, whose roots/rays can be exhibited as subsets of the roots/rays of E8. The rays of E7 are simply the 63 rays orthogonal to any ray of E8; these rays lie in a 7-dimensional space, and if one adjoins to them all their negatives, one gets the 126 roots of E7. The 63 rays of E7 form 135 bases, with each ray occurring in 15 bases. This system can be described by the symbol 63_{15} -135₇, and it is saturated. The basis table of E7 is easily extracted from that of E8 by picking out the 135 bases involving a particular ray and then dropping that ray from these bases. This construction shows that the symmetry group of E7 is a subgroup of index 240 of that of E8; thus its order is $192 \cdot 10!/240$, which agrees

with the figure of $8 \cdot 9!$ given by Coxeter for the order of the symmetry group of the associated 7-dimensional polytope 3_{21} . The rays of E7 can be used to give a proof of the KS theorem. Because the system is saturated, the proof requires showing that it is impossible to assign noncontextual 0/1 values to the rays in such a way that each of the 135 bases has just a single ray assigned the value 1 in it, and this is easily done using a “proof-tree” argument⁵. An alternative proof of the KS theorem based on the rays of E7 has been given by Ruuge[20].

If one picks the 36 rays orthogonal to any two nonorthogonal rays of E8, one gets the rays of the E6 system, and the roots of E6 are these 36 rays along with their negatives. The rays of E6 do not form even a single basis (i.e., a set of six mutually orthogonal rays) and so do not yield a proof of the KS theorem. This was pointed out by Ruuge[20], who discussed how two rotated copies of E6 could be superposed to give a system of rays that yields a proof of the KS theorem.

Another interesting subsystem of E8 is what we will term a Kernaghan-Peres (KP) set[13]. Such a set consists of 40 rays that form 25 bases, with each ray occurring in five bases, so that its symbol is 40_5 - 25_8 . Kernaghan and Peres[13] constructed such a set as the simultaneous eigenstates of five sets of mutually commuting observables of a system of three qubits. The caption to Fig.3 explains how KP sets can be extracted from the bases of E8 and the figure gives an example of a set constructed using this procedure. The caption to Fig.4 gives a simple procedure for obtaining all the parity proofs in a KP set[14] and the figure gives one example of each of the three types of parity proofs contained in the KP set of Fig.3 (see the beginning of Sec. 4 for an explanation of the notion of a parity proof). Finally, Fig.5 gives an example of a “pseudo” KP set that closely resembles a KP set but is not one, and in fact yields no proofs of the KS theorem.

⁵ The argument is a *reductio ad absurdum* one: one assumes that a noncontextual value assignment exists and then shows that it leads to a contradiction. Since E7 has a symmetry group that is transitive on its rays, one can begin, without loss of generality, by assigning an arbitrary ray the value 1. This forces all rays orthogonal to that ray to have the value 0, and one then finds that a basis with three rays having the value 0 appears. Assigning one of the remaining rays in this basis the value 1 forces a basis with five rays having the value 0 to appear. However assigning either of the remaining rays in this basis the value 1 forces a basis with all its rays assigned the value 0 to appear, which is not allowed. To avoid this conflict, one must proceed backwards along the chain and make alternative choices for the rays assigned the value 1 at every earlier step of the argument and see if any of these alternative possibilities leads to a valid value assignment. One then finds that all the alternatives lead to a situation in which at least one basis has all its rays assigned the value 0, showing that a valid value assignment does not exist and proving the KS theorem. The “proof-tree” leading to this contradiction has eight branches, with each branch leading to a contradiction at the fourth step.

1 7 62 66 70 73 107 111	1 64 94 7 66 62 34 19	101 102 70 111 73 88 85 107
29 115 33 11 74 61 5 52	52 1 107 61 73 7 33 29	5 11 66 70 74 62 111 115
64 34 85 102 19 88 101 94	62 92 116 1 47 42 70 107	78 110 7 111 66 58 112 73
58 78 92 42 47 110 112 116	66 38 1 10 23 73 103 70	56 7 111 53 37 113 62 107
10 23 38 103 37 56 53 113	29 115 47 78 58 33 11 116	112 52 42 74 61 92 5 110
	5 94 29 88 64 61 85 115	19 33 34 52 74 102 11 101
	52 5 11 29 23 38 37 56	53 113 74 115 10 33 61 103
	88 92 116 101 64 34 58 110	19 94 112 78 42 85 102 47
	94 34 103 23 88 37 53 102	10 64 19 38 113 56 101 85
	113 58 37 112 92 103 47 38	110 42 116 78 23 10 53 56

Fig. 3 A Kernaghan-Peres (KP) set can be extracted from the bases of E8 by choosing any five (“seed”) bases from one of the blocks in Fig.2 (or a block obtained from one of these blocks by applying a power of W to it) and supplementing them by 20 bases, chosen from the full set of 2025, that each have four rays from one seed basis and four rays from another. The 20 added bases always come in 10 complementary pairs, with the members of each pair originating in the same pair of seed bases and having no rays in common. In order that this construction gives rise to a KP set, it is necessary that each ray occurs once with 17 other rays and thrice with 6 other rays in the five bases in which it occurs. Shown above is a KP set constructed according to this procedure, with its five seed bases shown in the first column and its 10 pairs of complementary bases in the second and third columns; the seed bases were picked from the block in the top left corner of Fig.2 and each pair of complementary bases is shown on a line. It can be checked that this set of rays and bases has the symbol 40_5-25_8 and that each of its rays has the pattern of companions stated earlier.

1 64 94 7 66 62 34 19	1 64 94 7 66 62 34 19	1 64 94 7 66 62 34 19
52 1 107 61 73 7 33 29	52 1 107 61 73 7 33 29	52 1 107 61 73 7 33 29
62 92 116 1 47 42 70 107	62 92 116 1 47 42 70 107	62 92 116 1 47 42 70 107
66 38 1 10 23 73 103 70	66 38 1 10 23 73 103 70	56 7 111 53 37 113 62 107
29 115 47 78 58 33 11 116	29 115 47 78 58 33 11 116	29 115 47 78 58 33 11 116
5 94 29 88 64 61 85 115	5 94 29 88 64 61 85 115	5 94 29 88 64 61 85 115
52 5 11 29 23 38 37 56	52 5 11 29 23 38 37 56	53 113 74 115 10 33 61 103
88 92 116 101 64 34 58 110	88 92 116 101 64 34 58 110	88 92 116 101 64 34 58 110
94 34 103 23 88 37 53 102	94 34 103 23 88 37 53 102	94 34 103 23 88 37 53 102
110 42 116 78 23 10 53 56	113 58 37 112 92 103 47 38	113 58 37 112 92 103 47 38
64 34 85 102 19 88 101 94	64 34 85 102 19 88 101 94	1 7 62 66 70 73 107 111
	58 78 92 42 47 110 112 116	29 115 33 11 74 61 5 52
	10 23 38 103 37 56 53 113	64 34 85 102 19 88 101 94
		58 78 92 42 47 110 112 116
		10 23 38 103 37 56 53 113

Fig. 4 Parity proofs can be extracted from any KP set by picking one member from each of the 10 pairs of complementary bases and supplementing them with the needed seed bases. There are three types of parity proofs that can be constructed in this way, and they involve the addition of one, three or five seed bases. These proofs have the symbols $28_28_4-11_8$, $24_214_4-13_8$ and $20_220_4-15_8$, and one example of each, constructed from the KP set of Fig.3, is shown in the three columns above (with the seed bases always shown at the end). The 10 bases in the first step of the construction can be picked in $2^{10} = 1024$ ways, and all of them can be extended into valid parity proofs.

4 Parity proofs in the E8 system

We will say that a set of projectors in a Hilbert space of even dimension furnishes a parity proof of the KS theorem if the projectors form an odd number of bases in such a way that each projector occurs in an even number of the bases (a basis is any set of mutually orthogonal projectors that sums to the

1 7 62 66 70 73 107 111	1 64 94 7 66 62 34 19	101 102 70 111 73 88 85 107
64 34 85 102 19 88 101 94	26 76 9 119 73 70 66 1	7 111 54 89 77 107 62 17
17 89 9 76 54 26 119 77	28 50 71 39 107 1 7 73	66 93 6 70 36 99 111 62
99 71 36 28 93 39 6 50	18 83 1 111 65 62 73 8	66 40 70 97 107 98 7 68
83 8 40 97 98 18 65 68	89 102 19 9 76 17 88 64	26 101 119 85 34 77 94 54
	85 50 34 99 28 19 6 88	93 64 94 36 101 102 71 39
	101 68 34 18 8 88 97 64	19 102 94 85 98 83 40 65
	119 54 39 36 50 17 9 99	77 26 93 71 89 28 6 76
	54 76 40 8 26 97 83 17	9 98 18 77 68 65 89 119
	99 83 36 8 68 28 98 71	40 65 6 50 97 39 93 18

Fig. 5 The construction that gave rise to the KP set of Fig.3 also leads to many “pseudo” KP sets, like the one above. Although a 40₅-25₈ set, it is not a KP set because each of its rays occurs four times with a particular companion in five bases, which never happens in a KP set. Further, this “pseudo” KP set does not yield a proof of the KS theorem because it is possible to make valid noncontextual 0/1 assignments to its rays in many ways; one such assignment consists of assigning 1’s to the rays 1,85,17,93 and 68 and 0’s to all the others.

identity, and we will allow for the possibility that the projectors are not all of the same rank). Such a set of projectors proves the KS theorem because it is impossible to assign noncontextual 0/1 values to them in such a way that the sum of the values assigned to the projectors in any basis is always 1. Because an even-odd conflict makes this assignment impossible, we refer to this type of proof as a parity proof. In [3] it was pointed out that the E8 system has 2¹⁹⁴⁰ parity proofs in it, but no examples of such proofs were given. In this section we would like to describe a straightforward method we used to discover a large number of these proofs, and then present a few examples of them. These proofs are far more numerous and varied than those in the KP sets we know to be contained in the E8 system.

We will discuss only *critical* parity proofs, where by a critical proof we mean one that ceases to provide a proof of the KS theorem if even a single basis is dropped from it⁶. We restrict ourselves to critical proofs to avoid redundancy, since many noncritical proofs can often be reduced to the same critical proof. We present just a few of the more striking critical proofs we found from among the staggeringly large number that exist.

In Sec. 2 we introduced the letters A to H for each consecutive set of 15 rays of E8. These letters can be used to attach an 8-letter label to each basis. For example, the basis 1 7 62 66 70 73 107 111 would have the label AAEEEEHH. Viewed in terms of their labels (which specify how the rays of a basis are distributed over the triacontagons of E8), the bases fall into 33 families with distinct triacontagon profiles. We made the important discovery that if one looks at only the bases of a particular family, the parity proofs housed by them

⁶ Dropping a single basis from a parity proof leaves an even number of bases, which can never provide a parity proof of the KS theorem. However the reduced system may not admit a valid noncontextual value assignment to its rays, and so provide a proof of the KS theorem. If this happens, the original parity proof would not be deemed critical.

could be unearthed with relatively little effort. In the following subsections we discuss the parity proofs we found in a few of the families.

4.1 Type 1 Bases and their parity proofs

Let us term bases with the profile AAEEEEHH, BBFFFFGG, CCEEGGGG or DDDFFHHH as Type 1 bases. The 15 bases with each of these profiles give a parity proof that can be characterized by the symbol $15_430_2-15_8$. Figure 6 shows the proof given by the bases with profile AAEEEEHH (the proofs given by the other three profiles are very similar).

The Type 1 bases give rise to two other types of parity proofs if bases of different profiles can be combined. These proofs are characterized by the symbols $15_470_2-25_8$ and $45_450_2-35_8$ and there are 12 versions of each (all structurally identical, but involving different rays). The properties of the three different types of parity proofs made up only of Type 1 bases are summarized in the first row of Figure 7.

1 7 62 66 70 73 107 111	6 12 67 71 75 63 112 116	11 2 72 61 65 68 117 106
2 8 63 67 71 74 108 112	7 13 68 72 61 64 113 117	12 3 73 62 66 69 118 107
3 9 64 68 72 75 109 113	8 14 69 73 62 65 114 118	13 4 74 63 67 70 119 108
4 10 65 69 73 61 110 114	9 15 70 74 63 66 115 119	14 5 75 64 68 71 120 109
5 11 66 70 74 62 111 115	10 1 71 75 64 67 116 120	15 6 61 65 69 72 106 110

Fig. 6 A $15_430_2-15_8$ parity proof made up of 15 bases with profile AAEEEEHH. The rays 1-15 and 106-120 have multiplicity 2, while rays 61-75 have multiplicity 4.

4.2 Type 2 Bases and their parity proofs

We will term bases with the profile AABBEFF or CCDDGGHH as Type 2 bases. There are 30 bases with each of these profiles, and therefore 60 Type 2 bases in all. These bases contain just the two types of parity proofs shown in the second row of Figure 7.

Figure 8 shows two 36_2-9_8 proofs of this class that seem very similar at first sight, but are subtly different from one another. While both proofs involve 36 rays that occur two times each over 9 bases, the proof on the left always has the following pairs of rays occur together over the bases: (1,66), (7,62), (16,86), (22,87), (9,84), (19,64), (21,76), (14,89), (17,82), (2,72), (11,61), (27,77), (29,74), (26,81), (4,79), (12,67), (24,69) and (6,71). Thus each of these pairs of rays can be regarded as defining a two-dimensional subspace, or a rank-2 projector, and the proof can be reinterpreted as involving 18 rank-2 projectors that each occur twice over nine bases; this situation can be captured

Basis Type	Parity Proofs	Number
Type 1 (AAEEEEHH, BBFFFFGG, CCEEEEGG, DDFFHHHH)	15 ₄ 30 ₂ -15 ₈ 15 ₄ 70 ₂ -25 ₈ 45 ₄ 50 ₂ -35 ₈	4 12 12
Type 2 (AABBEEFF, CCDDGGHH)	36 ₂ -9 ₈ 60 ₂ -15 ₈	20 24
Type 3 (EEFFGGHH)	60 ₂ -15 ₈ , 52 ₂ 8 ₂ -17 ₈ , 16 ₄ 44 ₂ -19 ₈ , 1 ₆ 14 ₄ 45 ₂ -19 ₈ , 2 ₆ 12 ₄ 46 ₂ -19 ₈ 24 ₄ 36 ₂ -21 ₈ , 1 ₆ 22 ₄ 37 ₂ -21 ₈ , 2 ₆ 20 ₄ 38 ₂ -21 ₈ , 3 ₆ 18 ₄ 39 ₂ -21 ₈ , 4 ₆ 16 ₄ 40 ₂ -21 ₈ 6 ₆ 12 ₄ 42 ₂ -21 ₈ , 3 ₂ 428 ₂ -23 ₈ , 1 ₆ 30 ₄ 29 ₂ -23 ₈ , 2 ₆ 28 ₄ 30 ₂ -23 ₈ , 3 ₆ 26 ₄ 31 ₂ -23 ₈ 4 ₆ 24 ₄ 32 ₂ -23 ₈ , 5 ₆ 22 ₄ 33 ₂ -23 ₈ , 6 ₆ 20 ₄ 34 ₂ -23 ₈ , 7 ₆ 18 ₄ 35 ₂ -23 ₈ , 8 ₆ 16 ₄ 36 ₂ -23 ₈	700,326
Type 4 (AABBCCDD)	2 ₄ 32 ₂ -9 ₈ , 3 ₆ 2-9 ₈ , 8 ₄ 28 ₂ -11 ₈ , 7 ₄ 30 ₂ -11 ₈ , 13 ₄ 26 ₂ -13 ₈ 2 ₆ 15 ₄ 24 ₂ -15 ₈ , 1 ₈ 2 ₆ 9 ₄ 32 ₂ -15 ₈ , 1 ₈ 3 ₆ 11 ₄ 33 ₂ -17 ₈ , 12 ₆ 24 ₄ 24 ₂ -27 ₈	

Fig. 7 Parity proofs in E8. For each of the classes of bases in the first column, the second column lists the symbols of the parity proofs that exist and the third column the number of versions of each of the proofs. The listings for Type 1 and Type 2 bases are complete. For Type 3 bases, the listing in the second column is complete but only the total count of all the proofs has been included in the third column. For Type 4 proofs, only nine of the over hundreds of different types of proofs we found are listed. The left halves of the proof symbols in the second column indicate the numbers of rays of different multiplicities present in the proofs, with no attempt being made to associate the rays with projectors of different ranks. Figs. 8, 10 and 12 give examples of proofs in which this distinction is made.

in the symbol $18_2^2-9_4$, where the superscript on the left indicates the rank of the projectors and the subscript their multiplicity, and the subscript on the right that each of the 9 bases is made up of four rank-2 projectors. For the proof on the right, 18 of the rays can be paired into the rank-2 projectors (1,16), (62,87), (75,80), (70,90), (82,72), (65,85), (67,77), (11,26) and (6,21), while the remaining 18 rays are associated with rank-1 projectors; thus the symbol of this proof can be written as $9_2^2 18_1^1-9_6$, with the superscripts and subscripts on the left having the same meaning as before and the subscript on the right indicating that each basis is made up of six projectors (four of rank-1 and two of rank-2).

4.3 Type 3 Bases and their parity proofs

Type 3 bases are those with the profile EEFFGGHH. There are 45 such bases involving 60 rays, and they form a 60_6-45_8 system. Despite their small number, the bases of this system are a fecund lot and give rise to 20 different types of parity proofs, each of which can come in hundreds to thousands of versions. The symbols of all the possible proofs are shown in the third row of Figure 7. When the different versions of each of the proofs are taken into account, the total number of distinct proofs is 700,326. Figure 9 shows a

1 7 16 22 62 66 86 87	1 7 16 22 62 66 86 87
9 1 19 21 64 66 84 76	10 1 25 16 71 75 80 81
7 14 17 19 62 64 82 89	15 7 25 27 70 72 90 82
2 9 27 29 72 74 77 84	10 2 20 22 65 67 85 77
11 2 26 17 72 61 81 82	11 2 26 17 72 61 81 82
4 11 29 16 74 61 79 86	5 11 20 26 66 70 90 76
12 4 22 24 67 69 87 79	5 12 30 17 75 62 80 87
6 12 21 27 67 71 76 77	6 12 21 27 67 71 76 77
14 6 24 26 69 71 89 81	15 6 30 21 61 65 85 86

Fig. 8 Two parity proofs made up exclusively of Type 2 bases with the profile AABBEFF or CCDDGGHH. The proof on the left involves only rank-2 projectors and is characterized by the symbol $18_2^2-9_4$, while the proof on the right involves a mixture of rank-2 and rank-1 projectors and is characterized by the symbol $9_2^2 18_2^1-9_6$ (see text for explanation). If one ignores the distinction between rank-1 and rank-2 projectors and focuses only on the rays, both proofs can be described by the common symbol 36_2-9_8 (indicating that there are 36 rays that each occur twice over the nine bases).

61 63 81 88 105 94 114 115	68 70 88 80 97 101 106 107	67 70 82 85 98 99 119 108
62 64 82 89 91 95 115 116	69 71 89 81 98 102 107 108	65 72 90 77 103 94 118 109
63 65 83 90 92 96 116 117	72 74 77 84 101 105 110 111	66 73 76 78 104 95 119 110
64 66 84 76 93 97 117 118	73 75 78 85 102 91 111 112	69 61 79 81 92 98 107 113
65 67 85 77 94 98 118 119	62 65 77 80 93 94 114 118	71 63 81 83 94 100 109 115
66 68 86 78 95 99 119 120	63 66 78 81 94 95 115 119	74 66 84 86 97 103 112 118
67 69 87 79 96 100 120 106	66 69 81 84 97 98 118 107	75 67 85 87 98 104 113 119

Fig. 9 A $6_6 12_4 42_2-21_8$ parity proof made up of 21 Type 3 bases with the profile EEFGGHH. The rays of multiplicity 6 are 66,81,94,98,118,119, those of multiplicity 4 are 63,65,67,69,77,78,84,85,95,97,107,115, and all the other rays have multiplicity 2.

$6_6 12_4 42_2-21_8$ proof of this class involving rays of multiplicity 6,4 and 2.

4.4 Type 4 Bases and their parity proofs

Type 4 bases have the profile AABBCDD. There are 75 such bases involving 60 rays, and they form a $60_{10}-75_8$ system. We have found over 400 different types of parity proofs in this system, with each coming in anywhere from scores to thousands of versions. We show just nine of these proofs in the last row of Figure 7. There are no critical proofs with more than 27 bases in this class. The number of proofs in this class greatly exceeds those in the previous three classes combined. Figure 10 shows a 36_2-9_8 proof of this class and Fig. 11 shows a rather unusual proof consisting entirely of rank-2 projectors.

We end by presenting a proof of this class, in Fig.12, that involves 34 rays (2 of multiplicity four and 32 of multiplicity two) that occur over 9 bases. This proof is more economical than the best proofs found earlier in 8 dimensions, which involve 36 rays occurring an even number of times over 11 bases[13] or 81 bases[49] or 9 bases[14]. As explained in the caption to Fig.12, this proof can also be interpreted as involving 26 rank-1 projectors and 4 rank-2 projec-

1	7	19	25	33	34	48	52
7	13	25	16	39	40	54	58
10	1	28	19	42	43	57	46
3	10	18	16	40	37	46	49
5	12	20	18	42	39	48	51
11	3	26	24	33	45	54	57
4	11	26	28	34	37	58	55
13	5	20	22	43	31	52	49
12	4	22	24	45	31	51	55

Fig. 10 A 36_2-9_8 parity proof made up of 9 Type 4 bases with the profile AABBCDD. Twenty of the rays can be grouped into the rank-2 projectors (1,19), (7,25), (16,40), (10,46), (5,20), (12,51), (11,26), (22,31), (24,45) and (4,55), while the remaining rays define 16 rank-1 projectors. Also, eight of the bases involve 6 projectors (two of rank-2 and four of rank-1) while the remaining basis involves four rank-2 projectors. Thus a more descriptive symbol for this proof is $10_2^2 16_2^1-8_6 1_4$.

1	7	19	25	33	34	48	52
10	1	28	19	42	43	57	46
13	4	16	22	45	31	60	49
8	14	26	20	41	42	53	57
11	2	29	23	44	45	56	60
14	5	17	26	32	33	59	48
15	7	30	28	37	34	58	46
8	15	30	17	38	41	47	59
4	11	29	16	37	38	58	47
10	2	20	22	43	44	49	53
13	5	23	25	31	32	52	56

Fig. 11 A parity proof made up of 11 Type 4 bases with the profile AABBCDD. It consists of 44 rays that each occur twice over the 11 bases, so its symbol is 44_2-11_8 . However a more careful examination shows that the rays can be paired into the 22 rank-2 projectors (1,19), (2,44), (4,16), (5,32), (7,34), (25,52), (33,48), (17,59), (14,26), (15,30), (11,29), (37,58), (38,47), (8,41), (28,46), (23,56), (13,31), (45,60), (22,49), (10,43), (20,53) and (42,57) that each occur twice over the bases. Thus a more descriptive symbol for this proof would be $22_2^2-11_4$, with the subscript of 4 on the right indicating that there are four rank-2 projectors in each basis.

tors that occur over 9 bases.

5 Discussion

We pointed out at the end of Sec. 3 that the bases of E8 have 33 different triacontagon profiles. Our survey of parity proofs in Sec. 4 covered just four of these profiles, so it is clear that we have left the vast majority of the proofs untouched. The basis table of E8 presented in this paper serves a convenient template for displaying all the proofs in this gargantuan system.

It is interesting that the triacontagonal representations of both the 600-cell and Gosset's polytope lead to some of the simplest parity proofs contained in them. In the case of the 600-cell, the vertices project into four triacontagons, with two of the triacontagons uniting to yield a parity proof of 15 bases and the other two triacontagons yielding the complementary proof (i.e, one involving

1	7	19	25	33	34	48	52
4	10	22	28	36	37	51	55
7	13	25	16	39	40	54	58
10	1	28	19	42	43	57	46
3	10	18	16	40	37	46	49
5	12	20	18	42	39	48	51
12	4	27	25	34	31	55	58
3	10	25	27	33	36	57	54
13	5	20	22	43	31	52	49

Fig. 12 A $32_2 24_4$ -9₈ parity proof made up of 9 Type 4 bases with the profile AABBCDD, with rays 25 and 10 being of multiplicity four and all the others of multiplicity two. However the pairs of rays (1,19), (4,55), (5,20) and (16,40) always occur together over the bases, with each pair occurring twice. Interpreting these pairs as rank-2 projectors and the remaining 26 rays as rank-1 projectors allows us to attach the more descriptive symbol $4_2^2 2_4^1 24_2^1$ -871₈ to this proof, where the subscripts in the second half of the symbol indicate that there are eight bases of 7 projectors (with 6 being of rank-1 and 1 of rank-2) and one basis of 8 projectors (with all being of rank-1).

all the rays not present in the earlier proof). In the case of Gosset’s polytope, the vertices project into eight triacontagons, and one can construct a parity proof (actually four different proofs) by picking out 15 bases that each span all the triacontagons in the same way. The great virtue of the triacontagonal representation for Gosset’s polytope (or E8) is, of course, that it allows the bases to be organized into smaller families that are more easily searched for their parity proofs. Although we have unearthed only a tiny fraction of the parity proofs present in E8, their variety and intricacy seems to exceed that in any of the other systems we have studied to date. This is doubtless due to the large basis table of E8 (at 2025 bases, a record) and its huge symmetry group (of over 10^8 elements).

A comment should be made about the experimental measurements needed to realize the bases of E8, on which all the parity proofs of this paper depend. It might be asked if the projectors corresponding to some of the bases can be realized as simultaneous eigenstates of commuting three-qubit observables that are tensor products of Pauli operators of the individual qubits. While this is true of some of the bases, such as the ones we have identified as the Kernaghan-Peres sets, it is not true of the bases in general. The simplest way of generating an arbitrary basis from the computational basis is by following Eq.(1) and applying a product of the appropriate powers of the three unitary operators U , V and W . Designing efficient quantum gates for these operators is an interesting problem that we will not take up here. However it seems worth pursuing because a recent experiment[50] has successfully generated several KS sets in a three-qubit system and holds out the possibility of eventually generating the more complex sorts of KS sets considered here.

It was pointed out in [24] that any KS proof can be converted into an inequality that is satisfied by any noncontextual hidden variables theory but violated in measurements carried out on an arbitrary quantum state. It might

be asked what the extent of the violation is for the parity proofs discussed in this paper. The answer to this question has already been given in an earlier work of ours[18]. We showed there that for any basis-critical parity proof (i.e., one which fails if even a single basis is omitted from it), the upper bound of the inequality for any noncontextual hidden variable theory is $B - 2$ (where B is the number of bases in the proof) whereas quantum mechanics predicts the value of B . This gap of 2 between the values predicted by hidden variable theories and quantum mechanics is a universal feature of all basis critical parity proofs. Thus the present proofs do not offer any particular advantage, from this point of view, over the many similar proofs[12, 14, 16, 18, 19] we found earlier.

Gosset's polytope is the real representative of a complex polytope known as Witting's polytope[6]. Coxeter[51] has carried out a systematic study of a large number of complex polytopes. It is possible that the ray systems derived from some of them might yield new proofs of the KS theorem. Whether this is true, and of what use it might be, are matters that remain to be explored.

Acknowledgements. One of us (PKA) would like to thank David Richter for stimulating his interest in E8 and supplying him with a copy of Ref.[7], which planted the seed for this work.

References

1. A.Marrani and P.Truini: Exceptional Lie algebras, $SU(3)$ and Jordan pairs: part 2. Zorn-type representations. *J. Phys. A: Math. Theor.* **47** 265202 (30pp)(2014). This paper has references to many uses of E8 in theoretical physics.
2. E.P.Specker: The Logic of Propositions which are not Simultaneously Decidable. *Dialectica* **14**, 239-246 (1960); S. Kochen, E.P.Specker: The problem of hidden variables in quantum mechanics. *J. Math. Mech.* **17**, 59-88 (1967). See also J.S. Bell: On the problem of hidden variables in quantum mechanics. *Rev. Mod. Phys.* **38**, 447-452 (1966). Reprinted in J.S. Bell, *Speakable and Unspeakable in Quantum Mechanics*, (Cambridge University Press, Cambridge, 1987).
3. P.Lisoněk, R.Raussendorf and V.Singh: Generalized parity proofs of the Kochen-Specker theorem. arXiv:1401.3035v1 (2014).
4. A.Ruuge and F.van Oystaeyen: Saturated Kochen-Specker type configuration of 120 projective lines in eight-dimensional space and its group of symmetry. *J. Math. Phys.* **46**, 052109 (28pp) (2005).
5. H.S.M.Coxeter: *Regular Polytopes* Dover, New York (1973), Ch.11.
6. H.S.M.Coxeter and G.C.Shephard: Portraits of a family of complex polytopes. *Leonardo* **25**, 239-244 (1992).
7. D.Richter: Triacontagonal coordinates for the $E(8)$ root system. arXiv:0704.3091 (2007).
8. A. Peres: Two simple proofs of the Kochen-Specker theorem. *J. Phys. A* **24**, L175-L178 (1991).
9. M. Kernaghan: Bell-Kochen-Specker theorem for 20 vectors. *J. Phys. A* **27**, L829-L830 (1994).
10. A. Cabello, J.M. Estebaranz, G. García-Alcaine: Bell-Kochen-Specker theorem: A proof with 18 vectors. *Phys. Lett. A* **212**, 183-187 (1996).
11. M. Pavičić, N.D. Megill, J.P. Merlet: New Kochen-Specker sets in four dimensions. *Phys. Lett. A* **374**, 2122-2128 (2010); M. Pavičić, J.P. Merlet, B.D. McKay, N.D. Megill: Kochen-Specker vectors. *J. Phys. A* **38**, 1577-1592 (2005).
12. M.Waegell and P.K.Aravind: Parity proofs of the Kochen-Specker theorem based on 60 complex rays in four dimensions. *J. Phys. A: Math. Theor.* **44**, 505303 (15pp) (2011).
13. M. Kernaghan, A. Peres: Kochen-Specker theorem for eight-dimensional space. *Phys. Lett. A* **198**, 1-5 (1995).
14. M.Waegell, P.K.Aravind: Proofs of the Kochen-Specker theorem based on a system of three qubits. *J. Phys. A: Math. Theor.* **45**, 405301 (13pp) (2012).
15. D.P.DiVincenzo and A.Peres: Quantum code words contradict local realism. *Phys. Rev. A* **55**, 4089-4092 (1997).
16. M.Waegell, P.K.Aravind: Proofs of the Kochen-Specker theorem based on the N-qubit Pauli group. *Phys. Rev. A* **88**, 012102 (10pp) (2013).
17. M.Waegell and P.K.Aravind: Parity Proofs of the Kochen-Specker Theorem Based on the 24 Rays of Peres. *Found. Phys.* **41**, 1786-1799 (2011).
18. M.Waegell, P.K.Aravind, N.D.Megill, M. Pavičić: Parity Proofs of the Bell-Kochen-Specker Theorem Based on the 600-cell. *Found. Phys.* **41**, 883-904 (2011).
19. M.Waegell, P.K.Aravind: Parity Proofs of the Kochen-Specker theorem based on the 120-cell. *Found. Phys.* **44**, 1085-95 (2014).
20. A.Ruuge: Exceptional and non-crystallographic root systems and the Kochen-Specker theorem. *J. Phys. A: Math. Theor.* **40** 28492859 (2007).
21. P.Lisoněk, P.Badziąg, J.R.Portillo, A.Cabello: Kochen-Specker set with seven contexts. *Phys. Rev. A* **89** 042101 (6pp) (2014).
22. G.Canas, M.Arias, S.Etcheverry, E.S.Gómez, A.Cabello, G.B.Xavier, G.Lima: Applying the Simplest Kochen-Specker Set for Quantum Information Processing. *Phys. Rev. Lett.* **113** 090404 (2014).
23. A. Cabello: Experimentally Testable State-Independent Quantum Contextuality. *Phys. Rev. Lett.* **101**, 210401 (4pp) (2008).
24. P. Badziąg, I. Bengtsson, A. Cabello, I. Pitowsky: Universality of State-Independent Violation of Correlation Inequalities for Noncontextual Theories. *Phys. Rev. Lett.* **103** 050401 (4pp) (2009).

25. G. Kirchmair, F. Zähringer, R. Gerritsma, M. Kleinmann, O. Gühne, A. Cabello, R. Blatt, C.F. Roos: State-independent experimental test of quantum contextuality. *Nature* **460**, 494-497 (2009).
26. H. Bartosik, J. Klep, C. Schmitzer, S. Sponar, A. Cabello, H. Rauch, Y. Hasegawa: Experimental Test of Quantum Contextuality in Neutron Interferometry. *Phys. Rev. Lett.* **103**, 040403 (4pp) (2009).
27. E. Amselem, M. Rådmark, M. Bourennane, A. Cabello: State-Independent Quantum Contextuality with Single Photons. *Phys. Rev. Lett.* **103**, 160405 (4pp) (2009).
28. O. Moussa, C.A. Ryan, D.G. Cory, R. Laflamme: Testing Contextuality on Quantum Ensembles with One Clean Qubit. *Phys. Rev. Lett.* **104**, 160501 (4pp) (2010).
29. L. Aolita, R. Gallego, A. Acín, A. Chiuri, G. Vallone, P. Mataloni, A. Cabello: Fully nonlocal quantum correlations. *Phys. Rev.* **A85**, 032107 (8pp) (2012).
30. V. D'Ambrosio, I. Herbauts, E. Amselem, E. Nagali, M. Bourennane, A. Cabello: Experimental Implementation of a Kochen-Specker Set of Quantum Tests. *Phys. Rev. X* **3**, 011012 (10pp) (2009).
31. T.S. Cubitt, D. Leung, W. Matthews, A. Winter: Improving Zero-Error Classical Communication with Entanglement. *Phys. Rev. Lett.* **104**, 230503 (4pp) (2010).
32. D. Hu, W. Tang, M. Zhao, Q. Chen, S. Yu, and C. H. Oh: Graphical nonbinary quantum error-correcting codes. *Phys. Rev.* **A78**, 012306 (11pp) (2008).
33. R. Raussendorf and H. J. Briegel: A One-Way Quantum Computer. *Phys. Rev. Lett.* **86** 5188-5191 (2001).
34. S. Abramsky, in *In Search of Elegance in the Theory and Practice of Computation* Lecture Notes in Computer Science Volume 8000 (Springer Berlin Heidelberg), pp 13-35 (2013).
35. O. Gühne, C. Budroni, A. Cabello, M. Kleinmann, J.-Å. Larsson: Bounding the quantum dimension with contextuality. *Phys. Rev.* **A89**, 062107 (2014).
36. Y. Aharonov, F. Colombo, S. Popescu, I. Sabadini, D.C. Struppa, J. Tollaksen: The quantum pigeonhole principle and the nature of quantum correlations. arXiv: 1407.3194 (2014).
37. S. Yu and C.H. Oh: Quantum pigeonhole effect, Cheshire cat and contextuality: arXiv: 1408.2477 (2014).
38. A. Rae and T.Forgan: On the implications of the Quantum-Pigeonhole Effect: arXiv: 1412.1333 (2014).
39. D.A. Meyer: Finite Precision Measurement Nullifies the Kochen-Specker Theorem. *Phys. Rev. Lett.* **83**, 3751-3754 (1999).
40. A. Kent: Noncontextual Hidden Variables and Physical Measurements. *Phys. Rev. Lett.* **83**, 3755-3757 (1999).
41. J. Barrett and A. Kent: Non-contextuality, finite precision measurement and the Kochen-Specker theorem. *Stud. Hist. Philos. Mod. Phys.* **35**, 151-176 (2004).
42. N.D. Mermin: A Kochen-Specker Theorem for Imprecisely Specified Measurement. arXiv:9912081.
43. A.A. Klyachko, M.A. Can, S. Binicioğlu and A.S. Shumovsky: Simple Test for Hidden Variables in Spin-1 Systems. *Phys. Rev. Lett.* **101**, 020403 (2008).
44. Y.C. Liang, R.W. Spekkens, H.M. Wiseman: Specker's parable of the overprotective seer: A road to contextuality, nonlocality and complementarity. *Phys. Rep.* **506**, 1 (2011).
45. P. Badziąg, I. Bengtsson, A. Cabello, H.H. Granström and J.-Å. Larsson: Pentagrams and Paradoxes. *Found Phys* **41**, 414-423 (2011).
46. S.Yu and C.H.Oh: State-Independent Proof of Kochen-Specker Theorem with 13 Rays. *Phys. Rev. Lett.* **108**, 030402 (2012).
47. R.W.Spekkens: The Status of Determinism in Proofs of the Impossibility of a Noncontextual Model of Quantum Theory. *Found Phys* **44** 11251155 (2014).
48. M.Howard, J.Wallman, V.Veitch, J.Emerson: Contextuality supplies the 'magic' for quantum computation. *Nature* **510** 351-355 (2014).
49. A. Cabello, J.M. Estebaranz, G. García-Alcaine: Recursive proof of the Bell-Kochen-Specker theorem in any dimension $n > 3$. *Phys. Lett. A* **339**, 425-429 (2005).
50. G.Canas, S.Etcheverry, E.S.Gómez, C.Saavedra, G.B.Xavier, G.Lima, A.Cabello: Experimental implementation of an eight-dimensional Kochen-Specker set and observation of its connection with the Greenberger-Horne-Zeilinger theorem. *Phys. Rev.* **A90**, 012119 (2014).

51. H.S.M.Coxeter: *Regular Complex Polytopes* Cambridge University Press, New York (1991).

